

Math 114 Discrete Mathematics
First Midterm Answers
February 2018

Scale. 90–100 A, 79–89 B, 67–78 C. Median 88.

1. Negating propositions. [15; 5 points each part] For each of the following propositions, write the negation of the proposition so that negations only appear immediately preceding predicates; there should be no negations of conjunctions, disjunctions, or quantifiers.

a. $P(x) \wedge P(y)$

The negation $\neg(P(x) \wedge P(y))$ can be converted using De Morgan's laws to $\neg P(x) \vee \neg P(y)$.

b. $\forall x P(x)$

Passing the negation in $\neg \forall x P(x)$ past the quantifier changes it from a universal quantifier to an existential one: $\exists x \neg P(x)$.

c. $\forall x \exists y (P(x) \rightarrow Q(x, y))$

Passing the negation past the quantifiers in $\neg \forall x \exists y (P(x) \rightarrow Q(x, y))$ yields $\exists x \forall y \neg (P(x) \rightarrow Q(x, y))$. Then, the negation of the implication $(P(x) \rightarrow Q(x, y))$ gives $P(x) \wedge \neg Q(x, y)$. So the final answer is $\exists x \forall y (P(x) \wedge \neg Q(x, y))$.

2. On truth tables. [20; 10 points each part]

a. Use a truth table to determine whether $(p \vee q) \rightarrow r$ is logically equivalent to $(p \rightarrow r) \wedge (q \rightarrow r)$. Explain in a sentence why your truth table says whether they are logically equivalent or not.

p	q	r	$(p \wedge q) \rightarrow r$	$(p \rightarrow r) \wedge (q \rightarrow r)$
T	T	T	T	T
T	T	F	F	F
T	F	T	T	T
T	F	F	T	F
F	T	T	T	T
F	T	F	T	F
F	F	T	T	T
F	F	F	T	T

Since the columns for the two propositions are the same in all eight rows, therefore the propositions are logically equivalent.

b. Use a truth table to determine whether $(p \wedge r \rightarrow q) \wedge (q \rightarrow p) \rightarrow (p \vee \neg q \vee r)$ is a tautology, a contradiction, or a contingent proposition. Explain in a sentence why your truth table shows whether it is a tautology, a contradiction,

or a contingent proposition.

p	q	r	$(p \wedge r \rightarrow q)$	\wedge	$(q \rightarrow p)$	\rightarrow	$(p \vee \neg q \vee r)$
T	T	T	T	T	T	T	T
T	T	F	F	T	T	T	T
T	F	T	T	F	F	T	T
T	F	F	F	T	T	T	T
F	T	T	F	T	F	F	T
F	T	F	F	T	F	F	F
F	F	T	F	T	T	T	T
F	F	F	F	T	T	T	T

This is a tautology since it's true in all eight rows.

3. Interpretation of symbolic expressions. [25; 5 points each part] True/false.

a. $\forall x (x^2 \geq 0)$. *True.* The square of every real number is nonnegative.

b. $\exists y \forall x (x < y)$. *False.* There is no number y greater than all numbers. Infinity is not a real number.

c. $\exists x \exists y (y^2 - x^2 = 1)$. *True.* For example, $x = 0$ and $y = 1$.

d. $\forall x \exists y (y^2 - x^2 = 1)$. *True.* Take $y = \pm \sqrt{1 + x^2}$.

e. $\exists y \forall x (y^2 - x^2 = 1)$. *False.* Given y , there are at most two values of x that satisfy this equation, $x = \pm \sqrt{y^2 - 1}$.

4. On proofs. [15] Prove that for any positive integer n , if 3 divides n^2 , then 3 divides n . Here, "divides" means divides without remainder. [Suggestion: one way you can do this is by a proof by contradiction using cases. There are 3 cases. Case a: 3 divides n without remainder (which you're trying to show). Case b: there is a remainder of 1 when 3 divides n . Case c: there is a remainder of 2 when 3 divides n .]

The proof that follows is based on the suggestion. There are various ways it can be written up.

Proof: Suppose that 3 divides n^2 . One of the three cases a, b, and c described in the suggestion holds.

Case b: Suppose there is a remainder of 1 when 3 divides n . Then $n = 3k + 1$ for some integer k . Then $n^2 = (3k + 1)^2 = 9k^2 + 6k + 1$. Since 3 divides $9k^2 + 6k$, therefore when 3 divides n^2 there is a remainder of 1. So case b cannot occur when 3 divides n^2 .

Case c: Suppose there is a remainder of 2 when 3 divides n . Then $n = 3k + 2$ for some integer k . Then $n^2 = (3k + 2)^2 = 9k^2 + 12k + 4$. Since 3 divides $9k^2 + 12k + 3$, therefore when 3 divides n^2 there is a remainder of 1 again. So case c cannot occur when 3 divides n^2 .

That leaves only case a in which 3 divides n .

Therefore, if 3 divides n^2 , then 3 also divides n . Q.E.D.

There are also proofs that use other properties of positive integers such as the unique factorization theorem. We'll discuss that later in chapter 3.

5. On sets. [25] True/false.

a. If $|A| = |B|$, then $|A| = |B|$. (Recall that $|A|$ is the cardinality of the set A .) *False*. A counterexample is given by $A = \{1, 2\}$ and $B = \{3, 4\}$.

b. If $A = \{1, 3, 5, 7, 9\}$, then $|\mathcal{P}(A)| = 32$. (Recall that $\mathcal{P}(A)$ is the powerset of A .) *True*. $|\mathcal{P}(A)| = 2^{|A|} = 2^5 = 32$.

c. If $A \cap B = A$, then $A \cup B = B$. *True*. Both equations are equivalent to $A \subseteq B$.

d. $A \cap B \subseteq C$ implies $A \cup B \cup C \subseteq A \cup B$. *False*. One counterexample is given by $A = \{1\}$, $B = \{2\}$, $C = \{1, 2, 3\}$.

e. The composition of two one-to-one functions is also a one-to-one function. (Recall that a one-to-one function is also called an injection.) *True*.